

* Particle-Hole Picture

* For N -body systems, it is useful to switch to basis states labeled as deviations (i.e., excitations of) from some reference N -body Slater Det. $|\Phi\rangle$ rather than states w/ N - a^\dagger acting on $|0\rangle$ where all N -particles explicitly active




$$|\Phi\rangle = \prod_{i=1}^N a_i^\dagger |0\rangle$$


Important! Many different labeling conventions exist!
Here, I will stick to

a, b, c, \dots = particle states
 i, j, k, \dots = hole states
 r, s, t, \dots = unrestricted

1 particle-1 hole (ph) excitations! $|\Phi_i^a\rangle = a_a^\dagger a_i |\Phi\rangle$



2p-2h excitations! $|\Phi_{ij}^{ab}\rangle = a_a^\dagger a_i a_b^\dagger a_j |\Phi\rangle$
 $= a_a^\dagger a_b^\dagger a_j a_i |\Phi\rangle$



$$\Rightarrow \mathbb{1}_{\mathcal{F}_N} = |\Phi\rangle\langle\Phi| + \sum_{a,i} |\Phi_i^a\rangle\langle\Phi_i^a|$$

$$+ \sum_{\substack{a < b \\ i < j}} |\Phi_{ij}^{ab}\rangle\langle\Phi_{ij}^{ab}|$$

$$+ \dots + \sum_{\substack{a < b < \dots < l \\ i < j < \dots}} |\Phi_{ij\dots}^{ab\dots}\rangle\langle\Phi_{ij\dots}^{ab\dots}|$$

Re-define creation/annihilation operators $(a, a^\dagger) \rightarrow (b, b^\dagger)$

$$\begin{aligned} a_a^\dagger &= b_a^\dagger & a_i &= b_i^\dagger \\ a_a &= b_a & a_i^\dagger &= b_i \end{aligned}$$

Why is this useful? ① $\{a_r, a_s^\dagger\} = \delta_{rs} \longrightarrow \{b_r, b_s^\dagger\} = \delta_{rs}$

i.e., canonical transf.

$$\textcircled{2} b_r |\Phi\rangle = 0$$

$$\langle\Phi| b_r^\dagger = 0$$

$$\textcircled{3} |\Phi_i^a\rangle = a_a^\dagger a_i |\Phi\rangle = b_a^\dagger b_i^\dagger |\Phi\rangle$$

etc...

i.e., mathematically just like

$$|d_1 d_2 \dots d_N\rangle = a_1^\dagger a_2^\dagger \dots a_N^\dagger |0\rangle$$

$$a_\alpha |0\rangle = 0$$

etc.

Normal Order w/respect to $|\Phi\rangle$

$$\{B_1 B_2 \dots B_n\} \equiv (-1)^P b_{p_1}^\dagger \dots b_{p_n}$$

Notation: to distinguish N-ordering wrt. $|\Phi\rangle$ from N-ordering wrt $|0\rangle$, // **
I will use $\{ \dots \}$ instead of $N[\dots]$.

ex: $\{b_i b_j^\dagger\} = -b_j^\dagger b_i$

$$\{b_i b_a^\dagger b_j b_b^\dagger\} = b_a^\dagger b_j b_i b_b^\dagger = -b_a^\dagger b_j^\dagger b_b b_i \text{ etc}$$

Note: often, we don't even bother going thru the notational step of $(a, a^\dagger) \rightarrow (b, b^\dagger)$

eg, the previous 2 examples $\{b_i b_j^\dagger\} = \{a_i^\dagger a_j\} = -a_j a_i^\dagger$

$$\{b_i b_a^\dagger b_j b_b^\dagger\} = \{a_i^\dagger a_a a_j a_b^\dagger\} = a_a^\dagger a_j a_i^\dagger a_b$$

i.e., the curly brackets are sufficient context that we push all $a_i^\dagger + a_a$ to the right and all $a_i + a_a^\dagger$ to the left when N-ordering wrt $|\Phi\rangle$

Contractions wrt $|\Phi\rangle$: $\overline{B_q B_r} \equiv B_q B_r - \{B_q B_r\}$

$$\overline{b_q b_r} = \overline{b_q^\dagger b_r^\dagger} = \overline{b_q^\dagger b_r} = 0$$
$$\overline{b_q b_r^\dagger} = \delta_{qr}$$

$$\overline{B_q B_r} \equiv B_q B_r - \{B_q B_r\}$$

$$\overline{b_q b_r} = \overline{b_q^\dagger b_r^\dagger} = \overline{b_q^\dagger} b_r^\dagger = 0$$

$$\overline{b_q b_r^\dagger} = \delta_{qr}$$

* In terms of the a^\dagger, a -operators

$$\overline{b_q^\dagger b_r} = \delta_{qr} \Rightarrow \overline{a_a^\dagger a_b} = a_a^\dagger a_b - \{a_a^\dagger a_b\} = a_a^\dagger a_b + a_b^\dagger a_a = \delta_{ab}$$

$$\overline{a_i^\dagger a_j} = a_i^\dagger a_j - \{a_i^\dagger a_j\} = a_i^\dagger a_j + a_j^\dagger a_i = \delta_{ij}$$

we can write this more compactly w/ no restriction on the indices

$$\overline{a_q^\dagger a_r} = \delta_{qr} n_q$$

$$\overline{a_q a_r^\dagger} = \delta_{qr} \bar{n}_q$$

$$\overline{a_q a_r} = \overline{a_q^\dagger a_r^\dagger} = 0$$

$$\text{where } n_q = \begin{cases} 1 & \text{if } q \text{ occupied in } |\Phi\rangle \\ 0 & \text{else} \end{cases}$$

$$\bar{n}_q = 1 - n_q$$

Wick's theorem wrt $|\Phi\rangle$

$$A_1 \dots A_n = \{A_1 \dots A_n\}$$

$$+ \{ \overline{A_1 A_2} \dots A_n \} + \{ \overline{A_1 A_2} \dots \overline{A_1 A_n} \} + \text{all single contractions}$$

$$+ \{ \overline{A_1 A_2} \overline{A_3 A_4} \dots A_n \} + \text{all double contractions}$$

+ ...

$$+ \{ \overline{A_1 A_2} \overline{A_3 A_4} \dots \overline{A_1 A_n} \} + \text{all fully contracted}$$

$$\langle \Phi | A_1 \dots A_n | \Phi \rangle = \sum_{\text{fully contracted}}$$

Ex: find $\langle \Phi | \hat{H} | \Phi \rangle$

$$\begin{aligned} \langle \Phi | \hat{T} | \Phi \rangle &= \sum_{g,r} T_{gr} \langle \Phi | a_g^\dagger a_r | \Phi \rangle = \sum_{g,r} T_{gr} \langle \Phi | a_r^\dagger a_g | \Phi \rangle \\ &= \sum_{g,r} T_{gr} n_g \delta_{gr} \\ &= \sum_g T_{gg} n_g = \sum_{i=1}^N T_{ii} \end{aligned}$$

$$\begin{aligned} \langle \Phi | \hat{V} | \Phi \rangle &= \frac{1}{4} \sum_{qrst} \langle qr | v | st \rangle \left(\langle \Phi | \underbrace{a_q^\dagger a_r^\dagger a_s a_t}_{n_q n_r \delta_{rt} \delta_{qs}} | \Phi \rangle + \langle \Phi | \underbrace{a_q^\dagger a_r^\dagger a_s a_t}_{- \delta_{qt} \delta_{rs} n_q n_r} | \Phi \rangle \right) \\ &= \frac{1}{4} \sum_{qr} \left(\langle qr | v | qr \rangle - \langle qr | v | rq \rangle \right) n_q n_r \\ &\quad + \langle qr | v | qr \rangle \\ &= \frac{1}{2} \sum_{ij} \langle ij | v | ij \rangle \end{aligned}$$

Example: N-ordered H (wrt $|\Phi\rangle$)

$$\hat{H} = \sum_{gr} T_{gr} a_g^\dagger a_r + \frac{1}{4} \sum_{qrst} V_{qrst} a_q^\dagger a_r^\dagger a_s a_t$$

already in N-order wrt $|\Phi\rangle$. It is useful to put it in N-order wrt $|\Phi\rangle$.

* just apply Wick's theorem (wrt $|\Phi\rangle$)

$$T = \sum_{g,r} T_{gr} a_q^\dagger a_r = \sum_{g,r} T_{gr} (\{a_q^\dagger a_r\} + \overline{a_q^\dagger a_r}) = \sum_i T_{ii} + \sum_{g,r} T_{gr} \{a_q^\dagger a_r\}$$

$$\Rightarrow T \equiv \langle \Phi | T | \Phi \rangle + \hat{T}_N \quad (1)$$

$$\hat{V} = \frac{1}{4} \sum_{g,r,s,t} V_{grst} a_q^\dagger a_r^\dagger a_s a_t$$

$$= \frac{1}{4} \sum_{g,r,s,t} V_{grst} \left(\{g^\dagger r^\dagger t s\} + \{g^\dagger r^\dagger t s\} + \{g^\dagger r^\dagger t s\} \right.$$

$$+ \{g^\dagger r^\dagger t s\} + \{g^\dagger r^\dagger t s\}$$

$$\left. + \{g^\dagger r^\dagger t s\} + \{g^\dagger r^\dagger t s\} \right)$$

(here, N stands
for N-ordered
NOT # of
particles!)

$$\hat{V} = \langle \Phi | \hat{V} | \Phi \rangle + \frac{1}{2} \sum_{g,s} \{g^\dagger s\} \left(\sum_{i=1}^N V_{gisi} \right) + \frac{1}{4} \sum_{g,r,s,t} V_{grst} \{g^\dagger r^\dagger t s\} \quad (2)$$

Combining (1) + (2): $\hat{H} = E_{\text{ref}} + \sum_{g,r} f_{gr} \{a_q^\dagger a_r\} + \frac{1}{4} \sum_{g,r,s,t} V_{grst} \{a_q^\dagger a_r^\dagger a_s a_t\}$

$$= E_{\text{ref}} + \hat{F}_N + \hat{V}_N = E_{\text{ref}} + \hat{H}_N$$

$$E_{\text{ref}} = \langle \Phi | H | \Phi \rangle$$

$$f_{gr} = T_{gr} + \frac{1}{2} \sum_{i=1}^N V_{gisi}$$

* Especially useful as approximate treatment of 3N forces

Realistic nuclear hamiltonians have 3NF's

$$H = \sum_{q,r} T_{qr} a_q^\dagger a_r + \frac{1}{4} \sum V_{qrst} a_q^\dagger a_r^\dagger a_s a_t + \frac{1}{36} \sum W_{qrstuv} a_q^\dagger a_r^\dagger a_s^\dagger a_v a_u a_t$$

Complicated Calculations
(memory, more expensive operation)

* Can N-order H wrt $|\Phi\rangle$

$$H = E_{ref} + \hat{f}_N + \hat{\Gamma}_N + \hat{W}_N$$

0-body
(#)

1-body
 $\sum_{qr} f_{qr} \{a_q^\dagger a_r\}$

2-body
 $\frac{1}{4} \sum \Gamma_{qrst} \{a_q^\dagger a_r^\dagger a_s a_t\}$

3-body
 $\frac{1}{36} \sum W_{qrstuv} \{a_q^\dagger a_r^\dagger a_s^\dagger a_v a_u a_t\}$

Key point: Dominant parts of \hat{W} subsumed in E_{ref} , f_{qr} , & Γ_{qrst}

$$\text{e.g. } \Gamma_{qrst} = V_{qrst} + \sum_i W_{qristi}$$

$$f_{qr} = T_{qr} + \frac{1}{2} \sum_i V_{qirvi} + \frac{1}{4} \sum_{ij} W_{qijrvij}$$

\Rightarrow Excellent approximation $\hat{H} \approx E_{ref} + \hat{f}_N + \hat{\Gamma}_N$

no more expensive than "traditional" calculations that ignore 3N completely, yet it implicitly includes dominant 3N contributions

Derivation of HF equations

* Want to find best variational estimate of $g_0 E$ within the restricted class of trial wf's that are simple Slater determinants

$$E_{g_0} \leq E^{\text{HF}} = \text{Min}_{|\Phi\rangle \in \text{Slater det}} \frac{\langle \Phi | H | \Phi \rangle}{\langle \Phi | \Phi \rangle}$$

$$|\Phi\rangle = \prod_{i=1}^N a_i^\dagger |0\rangle \quad \text{the s.p. states } a_i^\dagger |0\rangle = |i\rangle \text{ are the a-priori unknown HF basis}$$

Reminder!: Use the convention $a, b, c, \dots = \text{particles}$
 $i, j, k, \dots = \text{holes}$
 $g, r, s, \dots = \text{unrestricted}$

* Expand HF basis in some fixed s.p. basis (e.g., HO basis)

$$|g\rangle = \sum_{\alpha} |\alpha\rangle \langle \alpha | g \rangle = \sum_{\alpha} |\alpha\rangle D_{\alpha g}$$

\uparrow HF basis \uparrow HO basis

$$\Rightarrow a_g^\dagger = \sum_{\alpha} C_{\alpha}^\dagger D_{\alpha g} \quad \sum_{\alpha} D_{\alpha g} D_{\alpha g'}^* = \delta_{g g'} \quad (D D^\dagger = D^\dagger D = 1)$$

$$a_g = \sum_{\alpha} C_{\alpha} D_{\alpha g}^*$$

$$C_{\alpha}^\dagger = \sum_g a_g^\dagger D_{\alpha g}^* \quad \text{and} \quad C_{\alpha} = \sum_g a_g D_{\alpha g}$$

* Recall, any unitary transf. amongst the N -lowest (i.e. hole states) orbitals
 Only changes $|\Phi\rangle$ by a phase

$$\begin{aligned}
 |\Phi'\rangle &= \prod_{i=1}^N \left(\sum_{j=1}^N U_{ij} a_j^\dagger \right) |0\rangle = \sum_{i_1} \cdots \sum_{i_N} U_{1i_1} U_{2i_2} \cdots U_{Ni_N} a_{i_1}^\dagger a_{i_2}^\dagger \cdots a_{i_N}^\dagger |0\rangle \\
 &= \text{Det } U \cdot |\Phi\rangle \\
 &\quad \text{irrelevant} \\
 &\quad \text{phase since } UU^\dagger = 1
 \end{aligned}$$

Therefore, there is no 1-to-1 correspondence between the basis $|i\rangle$
 & the S.D. $|\Phi\rangle$

* However, there is a 1-to-1 correspondence between $|\Phi\rangle$ and
 its 1-body density matrix

$$\begin{aligned}
 \rho_{\alpha\beta} &= \langle \Phi | c_\beta^\dagger c_\alpha | \Phi \rangle \\
 &= \sum_q \sum_r D_{\beta q}^* D_{\alpha r} \underbrace{\langle \Phi | a_q^\dagger a_r | \Phi \rangle}_{\delta_{qr} N_r} \\
 &= \sum_q D_{\beta q}^* D_{\alpha q} N_q = \sum_i D_{\beta i}^* D_{\alpha i}
 \end{aligned}$$

$$\Rightarrow \rho_{\alpha\beta} = \langle \Phi | c_\beta^\dagger c_\alpha | \Phi \rangle = \sum_{i=1}^N \langle \alpha | i \rangle \langle i | \beta \rangle$$

$$\Rightarrow \rho_{\alpha\beta} = \langle \Phi | c_{\beta}^{\dagger} c_{\alpha} | \Phi \rangle = \sum_{i=1}^N \langle \alpha | i \rangle \langle i | \beta \rangle$$

$$\Rightarrow \text{equivalently, } \rho_{\alpha\beta} = \langle \alpha | \hat{P} | \beta \rangle \quad \hat{P} = \sum_{i=1}^N |i\rangle \langle i|$$

and

$\hat{P}^2 = \hat{P}$ projector in Sp space onto
the subspace spanned by
occupied orbitals $|i\rangle$

* As we already showed, there is 1-to-1 correspondence between
a Slater $|\Phi\rangle$ + its \hat{P} -operator

* Also, recall that all Slater det. have $\hat{P}^2 = \hat{P}$



$$E_{\text{gs}} \leq E^{\text{HF}} = \text{Min}_{|\Phi\rangle \in \text{Slater}} E[|\Phi\rangle] \equiv \text{Min}_{|\Phi\rangle \in \text{Slater det}} \langle \Phi | H | \Phi \rangle$$

doing this variation
over all Slater det.

equivalent to

varying \hat{P} subject to

constraint $\hat{P}^2 = \hat{P}$

Express $E[\Phi] = \langle \Phi | H | \Phi \rangle$ in terms of 1-body density matrix

$$E[\Phi] = \sum_{\alpha\beta} T_{\alpha\beta} \underbrace{\langle \Phi | c_{\alpha}^{\dagger} c_{\beta} | \Phi \rangle}_{\textcircled{A}} + \frac{1}{4} \sum_{\alpha\beta\gamma\delta} V_{\alpha\beta\gamma\delta} \underbrace{\langle \Phi | c_{\alpha}^{\dagger} c_{\beta}^{\dagger} c_{\gamma} c_{\delta} | \Phi \rangle}_{\textcircled{B}}$$

$$\textcircled{A} = \langle \Phi | c_{\alpha}^{\dagger} c_{\beta} | \Phi \rangle = \langle \Phi | \overbrace{c_{\alpha}^{\dagger} c_{\beta}} | \Phi \rangle = P_{\beta\alpha}$$

$$\textcircled{B} = \underbrace{\langle \Phi | c_{\alpha}^{\dagger} c_{\beta}^{\dagger} c_{\gamma} c_{\delta} | \Phi \rangle}_{+ P_{\delta\beta} P_{\gamma\alpha}} + \underbrace{\langle \Phi | c_{\alpha}^{\dagger} c_{\beta}^{\dagger} c_{\delta} c_{\gamma} | \Phi \rangle}_{- P_{\gamma\beta} P_{\delta\alpha}}$$

$$\Rightarrow E[\rho] = \sum_{\alpha\beta} T_{\alpha\beta} P_{\beta\alpha} + \frac{1}{4} \sum_{\alpha\beta\gamma\delta} V_{\alpha\beta\gamma\delta} (P_{\delta\beta} P_{\gamma\alpha} - P_{\gamma\beta} P_{\delta\alpha})$$

re-label dummy indices

$$E[\rho] = \sum_{\alpha\beta} T_{\alpha\beta} P_{\beta\alpha} + \frac{1}{2} \sum_{\alpha\beta\gamma\delta} V_{\alpha\beta\gamma\delta} P_{\delta\beta} P_{\gamma\alpha}$$

Variation of the $E[\hat{P}]$

$$\delta E = E[\hat{P} + \delta \hat{P}] - E[\hat{P}]$$

// What kind of Variations $\delta \hat{P}$ can we make to stay in Slater determinant space for $|\Phi\rangle$?

(restricting to Slater det. $|\Phi\rangle \iff (\hat{P} + \delta \hat{P})^2 = (\hat{P} + \delta \hat{P})$ [ie, has to remain projector]
as we search over \hat{P}]

$$\Rightarrow (\hat{P} + \delta \hat{P})^2 = \hat{P}^2 + \hat{P} \delta \hat{P} + \delta \hat{P} \hat{P} + \mathcal{O}(\delta \hat{P}^2)$$

$$\Rightarrow (\hat{P} + \delta \hat{P})^2 = \cancel{\hat{P}} + \hat{P} \delta \hat{P} + \delta \hat{P} \hat{P} = \cancel{\hat{P}} + \delta \hat{P}$$

$$\boxed{\delta \hat{P} \hat{P} = (1 - \hat{P}) \delta \hat{P}} \quad \otimes$$

Now, let $(1 - \hat{P}) \equiv \hat{\sigma}$ ($\hat{P} + \hat{\sigma} = 1$, $\hat{P}^2 = \hat{P}$, $\hat{\sigma}^2 = \hat{\sigma}$, $\hat{P} \hat{\sigma} = 0$)

M. \otimes by \hat{P} from left: $\hat{P} \delta \hat{P} \hat{P} = \hat{P} \cancel{\hat{\sigma}} \delta \hat{P}$

M. \otimes by $\hat{\sigma}$ from right: $\delta \hat{P} \hat{P} \hat{\sigma} = \hat{\sigma} \delta \hat{P} \hat{\sigma}$

Thus we find that $\hat{P} \delta \hat{P} \hat{P} = \hat{\sigma} \delta \hat{P} \hat{\sigma} = 0$

$$\Rightarrow \boxed{\delta P_{ab} = \delta P_{ij} = 0 \quad (\text{hh + pp Variations Vanish})}$$

\Rightarrow Only variations of the ph sector $\delta P_{ai}, \delta P_{ia} \neq 0$

$$\text{So then, } \delta E = E[P+\delta P] - E[P] = \sum_{\alpha\beta} \frac{\delta E}{\delta P_{\alpha\beta}} \delta P_{\alpha\beta} = 0$$

$$\frac{\delta E}{\delta P_{\alpha\beta}} = \frac{\delta}{\delta P_{\alpha\beta}} \left[\sum_{\sigma\tau} T_{\sigma\tau} P_{\sigma\tau} + \frac{1}{2} \sum_{m\nu\sigma\tau} V_{m\nu\sigma\tau} P_{\nu\tau} P_{m\sigma} \right]$$

$$= T_{\alpha\beta} + \sum_{m\sigma} V_{m\alpha\sigma\beta} P_{m\sigma} \equiv h_{\alpha\beta}^{\text{HF}}$$

$$\delta E = 0 = \sum_{\alpha\beta} h_{\alpha\beta}^{\text{HF}} \delta P_{\alpha\beta} \quad \left(\text{basis indep, so write it in HF basis} \right. \\ \left. \text{where we know only } \delta P_{ai}, \delta P_{ia} \text{ non-zero} \right)$$

$$= \sum_{gr} h_{gr}^{\text{HF}} \delta P_{gr}$$

$$= \sum_{ai} h_{ai}^{\text{HF}} \delta P_{ai} = 0$$

Since δP_{ai} arbitrary, this implies

$$h_{ai}^{\text{HF}} = 0 = h_{ia}^{\text{HF}} \quad \left(\text{particle/hole states not mixed} \right)$$

$$h_{ai}^{\text{HF}} = T_{ai} + \sum_{j=1}^N V_{ajij} = 0$$

* recalling that $\hat{P} = \sum_{i=1}^N |i\rangle\langle i|$

$$\hat{\sigma} = \sum_{a=N+1}^{\infty} |a\rangle\langle a| = 1 - \hat{P}$$

$$\Rightarrow \hat{P} h^{\text{HF}} (1 - \hat{P}) = 0$$

$$\Rightarrow \hat{P} h^{\text{HF}} - h^{\text{HF}} \hat{P} = 0 \quad \hat{P} + h^{\text{HF}} \text{ simultaneously diagonalizable}$$

$$h^{\text{HF}} = \left[\begin{array}{c|c} \hat{P} h^{\text{HF}} \hat{P} & 0 \\ \hline 0 & \hat{\sigma} h^{\text{HF}} \hat{\sigma} \end{array} \right]$$

particle + hole blocks
can be diagonalized
separately

\Rightarrow Go ahead and diagonalize $\hat{P} + h^{\text{HF}}$ simultaneously

$$\begin{aligned} \hat{P} |q\rangle &= n_q |q\rangle & n_q &= \begin{cases} 1 & q < N \\ 0 & q > N \end{cases} \\ h^{\text{HF}} |q\rangle &= \epsilon_q |q\rangle \end{aligned}$$

$$h^{HF} |q\rangle = \epsilon_q |q\rangle \quad (*)$$

* Now expand unknown $|q\rangle = \sum_{\alpha} |\alpha\rangle D_{\alpha q}$

↑
Some known fixed basis (eg - H0)

* Then eq. (*) becomes

$$\sum_{\beta} \langle \alpha | h^{HF} | \beta \rangle D_{\beta q} = \epsilon_q D_{\alpha q}$$

$$\text{where } \langle \alpha | h^{HF} | \beta \rangle = \langle \alpha | T | \beta \rangle + \frac{1}{2} \sum_{\mu\nu} \langle \alpha | \mu | V | \beta \nu \rangle P_{\mu\nu}$$

Requires iterative soln since h^{HF} depends on the eigenfunctions $D_{\beta q}$

Via

$$P_{\mu\nu} = \sum_{i=1}^N \langle \mu | i \rangle \langle i | \nu \rangle = \sum_{i=1}^N D_{\mu i} D_{\nu i}^*$$

Schematic Soln Strategy

- ① Guess initial $P_{\mu\nu}^{(0)}$ (eg, take the fixed $|\alpha\rangle$ basis as the initial guess for HF states $\Rightarrow P_{\mu\nu}^{(0)} = \delta_{\mu\nu} N_{\mu}$)

iterate
fill
things
don't
change

② Build $\langle \alpha | h^{HF} | \beta \rangle$

③ diagonalize $\Rightarrow D_{\alpha q}^{(1)}$

$$+ \text{build } P_{\mu\nu}^{(1)} = \sum_i D_{\mu i}^{(1)} D_{\nu i}^{(1)*}$$

$$+ h_{\alpha\beta}^{(1)}$$

$$\text{Total } E_{\text{HF}} := \langle \Phi | H | \Phi \rangle$$

$$E^{\text{HF}} = \sum_{\lambda} T_{\lambda\lambda} + \frac{1}{2} \sum_{\lambda j} \langle \lambda j | V | \lambda j \rangle$$

$$* \text{ but } \langle i | h^{\text{HF}} | i \rangle = \varepsilon_i = T_{ii} + \sum_j \langle \lambda j | V | \lambda j \rangle$$

⇓

$$\begin{aligned} E^{\text{HF}} &= \sum_{\lambda=1}^N \varepsilon_{\lambda} - \frac{1}{2} \sum_{\lambda j} \langle \lambda j | V | \lambda j \rangle \\ &= \frac{1}{2} \left(\sum_{\lambda} (T_{\lambda\lambda} + \varepsilon_{\lambda}) \right) \end{aligned}$$

$|i\rangle$ are the converged,
Self consistent orbitals
from $h^{\text{HF}} |i\rangle = \varepsilon_i |i\rangle$

How it looks in $|\vec{r}, \sigma\rangle$ basis

$$h_{\alpha\beta} = t_{\alpha\beta} + \frac{1}{2} \sum_{m\nu} \langle \alpha m | V | \beta \nu \rangle \rho_{m\nu}$$

here, $|\alpha\rangle \rightarrow |\vec{r}, \sigma\rangle$

$$\rho_{m\nu} \rightarrow \rho(\vec{r}\sigma, \vec{r}'\sigma') = \sum_{i=1}^A \langle \vec{r}\sigma | i \rangle \langle i | \vec{r}'\sigma' \rangle$$

HF basis

* For a local, spin-indep. V : $(\vec{r}_1\sigma_1 \vec{r}_2\sigma_2 | V | \vec{r}_3\sigma_3 \vec{r}_4\sigma_4)$

} Some algebra

$$\delta(\vec{r}_1 - \vec{r}_2) \delta_{\sigma_1\sigma_3} \delta_{\sigma_2\sigma_4} \delta(\vec{r}_1 - \vec{r}_3) \delta(\vec{r}_2 - \vec{r}_4)$$