

The r-process of nucleosynthesis: overview of r-process sites




Gail McLaughlin

North Carolina State University

Possible astrophysical sites of the r-process

The r -process elements

e. g. Uranium-238 $Z=92$, $N=146 \rightarrow$ need lots of neutrons



rapid neutron capture as compared with beta decay

Whats the most important criteria you are looking for?

Whats the most important criteria you are looking for?

What astrophysical sites have a lot of neutrons *and* eject material?

How do you get neutrons?

How do you get neutrons?

1. They already exist and just need to be liberated
 - in nuclei
 - in neutron stars
2. You make them through the weak interactions, i.e. conversion of protons into neutrons

How do you judge a site?

How do you judge a site?

- plenty of neutrons
- can populate halo stars
- how often does it occur
- does it match the abundance pattern


Of course, there could be more than one site...

Observational r-process data


Observational Halo Stars:
two r-process sites

Figure from Cowan and Sneden (2004)

main r-process and weak
r-process or multiple weak



Solar Abundances



What would be your first guess?

- Neutrino driven wind of the supernovae
- Jets from core collapse supernovae
- Accretion disks from core collapse supernovae
- ONeMg supernovae
- low entropy outflows from supernovae
- He Shell of core collapse supernovae
- Supernova with sterile neutrinos
- Tidal ejection of neutron rich matter in neutron star mergers
- shocked ejecta from merger
- accretion disk outflows from mergers

Possible astrophysical sites of the r-process

- Neutrino Driven Wind of the Supernovae
- Jets from Core Collapse Supernovae
- Accretion Disks from Core Collapse Supernova
- ONeMg Supernovae
- low energy outflows from supernovae
- Supernova with sterile neutrinos
- Tidal ejection of neutron rich matter in neutron star mergers
- shocked ejecta from merger
- accretion disk outflows from mergers

Compact object mergers




figure from Korobkin 2012




figure from Surman 2008

Mergers have many signals

- Gravitation wave signal, primary target of next generation detectors
- Prime candidate for short duration gamma ray bursts
- Huge emission of neutrinos, but hard to detect
- optical signal powered by radioactive decay of newly formed elements
- chemical evolution, elements produced in mergers, later observed in stars

Interesting from a nucleosynthetic point of view, but also for many other reasons

Evolution of neutron star merger

- Inspiral driven by gravitational wave emission
- Until last moments of inspiral, neutron stars may essentially be treated as cold neutron stars
- merger results in formation of a shocked extremely rapidly spinning hypermassive neutron star
- later formation of a disk around a black hole
- Models under development!

Types of mass ejection

- Dynamical ejection
 - material tidally ejected from tails
 - matter ejected through collisional region
- Winds
 - accretion disk
 - hypermassive neutron star
- Outflows from viscous heating

What happens to all this ejecta from a nucleosynthesis perspective?

Electron Fraction

In order to get the r-process nuclei, prefer a lot of neutrons


$$Y_e = \frac{n}{p + n} \quad (1)$$

Want this to be low.

neutron stars start with low Y_e .

Of the types of outflow we have considered (dynamical, wind, viscous heating driven), which has lowest Y_e ?

Dynamically ejected material from newtonian calculation



Goriely et al 2011

Y_e is so low you could have fission cycling!

Why fission cycling is a good thing

Basic observation



Halo star data suggest that abundance pattern in 2nd & 3rd peak region is “robust”. Abundance pattern below 2nd peak shows variations between different stars.

Need robust mechanism for populating 2nd & 3rd peaks.

Fission Cycling?

Note: Data show rare earth/3rd peak stable, few data in 2nd peak region. Generally assumed that 2nd/3rd also stable.


Fission Cycling in the r-process



abundance in 3rd/2nd peak as
a fct of decreasing Y_e

Very little data on the relevant fission rates and daughter products



Dynamically Ejected Material from Newtonian Calculation



Goriely et al 2011

Where is the evidence that there is fission cycling going on?



What about calculations where neutrinos are included?



What has happened to the Y_e ? Why?

Fig. from Wanajo et al 2014

The abundance pattern when neutrinos are included



Have these elements been fission cycled? Why or why not?

How much stuff?

Estimates depend on the hydrodynamics & thermodynamics & neutrino transport. Recent estimates:

- winds: $\sim 2 \times 10^{-3} M_{\odot}$ Wanajo and Janka 2011
- tidal tail ejection: 10^{-2} to $10^{-3} M_{\odot}$ Goriely et al 2011, Korobkin et al 2012

Need to make $\sim 10^{-2} M_{\odot}$ to account for all r-process material in Galaxy.

Does it match the halo stars?

Unresolved issues:

Mergers evolve slowly, $\tau_{\text{coales}} \approx 10^{6-8}$ years. Not clear how to populate halo stars

Mergers are rare, suggesting there should be more scatter in the amount of r-process material in halo stars than is seen.

Does it match the halo stars?

